
GPR58 Rabbit pAb

Catalog Number: bs-13541R

Target Protein: GPR58

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Mouse (predicted:Human, Rat, Rabbit, Pig, Sheep, Cow)

Predicted MW: 40 kDa

Entrez Gene: 9287

Swiss Prot: Q9P1P5

Source: KLH conjugated synthetic peptide derived from human GPR58: 61-160/351.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Trace amines are endogenous molecules structurally related to classical biogenic amines that are linked to psychiatric conditions. A family of G-protein coupled receptors referred to as trace-amine-associated receptors (TAAR) are activated by trace amines and are present in very low levels in mammalian tissue. TAARs contain several structural features that are similar to the rhodopsin beta-adrenergic receptor superfamily, including the positions of the seven transmembrane regions that provide common ligand-binding pockets as well as the short N- and C-terminal domains. TAAR proteins are potential targets for drugs of abuse, such as amphetamine and MDMA, as well as neuropsychiatric disorders including schizophrenia, depression, and attention deficit disorder. TAAR-1 is a 340 amino acid protein that increases intracellular cAMP accumulation in response to beta-phenylethylamine and tyramine. TAAR-1 is associated with the detection of social cues, illustrating its significance as a therapeutic target.

VALIDATION IMAGES



Sample: Cerebellum (Mouse) Lysate at 40 ug Primary: Anti-GPR58(bs-10196R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 40 kD Observed band size: 50 kD



Sample: placenta (Mouse) Lysate at 40 ug Cerebrum (Mouse) Lysate at 40 ug Primary: Anti-GPR58(bs-13541R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 40 kD Observed band size: 50 kD