

**bs-20814R**

**[ Primary Antibody ]**

## SLC22A1 Rabbit pAb

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000)
<b>Clonality:</b> Polyclonal		<b>Reactivity:</b> Human, Mouse
<b>GeneID:</b> 20517	<b>SWISS:</b> O08966	
<b>Target:</b> SLC22A1		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from mouse SLC22A1: 51-150/556.		<b>Predicted MW.:</b> 61 kDa
<b>Purification:</b> affinity purified by Protein A		<b>Subcellular Location:</b> Cell membrane
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The protein encoded by this gene is involved in the sodium-independent transport and excretion of organic anions, some of which are potentially toxic. The encoded protein is an integral membrane protein and is found mainly in the kidney and in the placenta, where it may act to prevent potentially harmful organic anions from reaching the fetus. [provided by RefSeq, Jul 2008]		

### — VALIDATION IMAGES —



Sample: Stomach (Mouse) Lysate at 40 ug  
Primary: Anti-SLC22A1(bs-20814R) at 1/300  
dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 61kD Observed band size: 70kD



Sample: HepG2 Cell (Human) Lysate at 40 ug  
Primary: Anti-SLC22A1(bs-20814R) at 1/300  
dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 61kD Observed band size: 70kD