

bs-20292R**[Primary Antibody]****TSPAN6 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

DATASHEET

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Mouse (predicted: Human, Rat, Rabbit, Pig, Sheep, Dog, Horse) Predicted MW.: 28 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 7105	SWISS: O43657	
Target: TSPAN6		
Immunogen: KLH conjugated synthetic peptide derived from human TSPAN6: 1-100/245. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Tetraspanins are a group of hydrophobic membrane proteins that interact with a wide variety of proteins including intracellular signaling molecules, integrins and membrane receptors. Members of the tetraspanin family are characterized by the presence of four hydrophobic domains and play a role in cell development, activation, growth and motility. TSPAN6 (tetraspanin-6), also known as A15 homolog, putative NF-kappa-B-activating protein 321, T245 protein, tetraspanin TM4-D or TM4SF6 (transmembrane 4 superfamily member 6), is a 245 amino acid multi-pass membrane protein that belongs to the tetraspanin (TM4SF) family. TSPAN6 is expressed at high levels in brain, with lower levels of expression found in colon, lung, pancreas, prostate, retina and melanocytes, as well as fetal heart, liver and spleen. The gene encoding TSPAN6 maps to the human X chromosome.		

VALIDATION IMAGES

Paraformaldehyde-fixed, paraffin embedded (mouse heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TSPAN6) Polyclonal Antibody, Unconjugated (bs-20292R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.