

bs-25128R**[Primary Antibody]****GGNBP2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:400-800) IF (1:100-500) Reactivity: Mouse, Rat (predicted: Human, Pig, Sheep, Horse) Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 79893	SWISS: Q9H3C7	
Target: GGNBP2		
Immunogen: KLH conjugated synthetic peptide derived from human GGNBP2: 501-600/697.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: LCRG1 is a 697 amino acid protein that may be involved in spermatogenesis. LCRG1 interacts with gametogenetin, a protein primarily expressed in testis and ovary. Localized in the cytoplasm, LCRG associates with vesicular structures. LCRG1 is expressed in brain, placenta, lung, liver and kidney, with highest levels found in heart, pancreas and skeletal muscle. LCRG1 is strongly down- regulated in 40% of primary laryngeal carcinoma and in 6 of 10 various cancer cell lines. Phosphorylated on Serine 360, LCRG1 exists as three isoforms produced by alternative splicing events.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GGNBP2) Polyclonal Antibody, Unconjugated (bs-25128R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GGNBP2) Polyclonal Antibody, Unconjugated (bs-25128R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.