

GGNBP2 Rabbit pAb

Catalog Number: bs-25129R

Target Protein: GGNBP2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:400-800), IF (1:100-500)

Reactivity: Human, Mouse (predicted:Rat, Rabbit, Pig, Sheep, Cow, Horse)

Entrez Gene: 79893

Swiss Prot: Q9H3C7

Source: KLH conjugated synthetic peptide derived from human GGNBP2: 601-697/697.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: LCRG1 is a 697 amino acid protein that may be involved in spermatogenesis. LCRG1 interacts with gametogenetin, a protein primarily expressed in testis and ovary. Localized in the cytoplasm, LCRG associates with vesicular structures. LCRG1 is expressed in brain, placenta, lung, liver and kidney, with highest levels found in heart, pancreas and skeletal muscle. LCRG1 is strongly down-regulated in 40% of primary laryngeal carcinoma and in 6 of 10 various cancer cell lines. Phosphorylated on Serine 360, LCRG1 exists as three isoforms produced by alternative splicing events.


VALIDATION IMAGES




Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GGNBP2) Polyclonal Antibody, Unconjugated (bs-25129R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GGNBP2) Polyclonal Antibody, Unconjugated (bs-25129R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GGNBP2) Polyclonal Antibody, Unconjugated (bs-25129R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GGNBP2) Polyclonal Antibody, Unconjugated (bs-25129R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.