bs-0296G-BF750

## [ Secondary Antibodies ]

## Goat Anti-Mouse IgG H&L, BF750 conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Goat Clonality: Polyclonal	<b>lsotype:</b> IgG	Applications: IF (1:100-1000) Flow-Cyt (1:100-1000)
Target: Goat Anti-Mouse IgG H&L Purification: affinity purified by Protein G Concentration: 1.0 mg/ml		ICC/IF (1:100-1000) Excitation Spectrum: 750nm Emission spectrum: 780nm Reactivity: Mouse
glycerol.	with 1% BSA, 0.03% Proclin300 and 5 e year. Avoid repeated freeze/thaw cy	
serum with normal l is important for our molecules are produ adaptive immune re functions; to bind to recruit other cells an variability of the IgG	gG), is one of the most abundant prot evels between 8-17 mg/mL in adult bl defence against microorganisms and ced by B lymphocytes as a part of our sponse. The IgG molecule has two sep the pathogen that elicited the respon d molecules to destroy the antigen. T pool is generated by somatic recomb pecificities in an individual at a given to be 1011 variants.	ood. IgG the parate ise and to he ination