bs-25219R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

NALP4 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 147945 SWISS: Q96MN2

Target: NALP4

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a member of the nucleotide-

binding and leucine-rich repeat receptor (NLR) family, and is predicted to contain an N-terminal pyrin effector domain (PYD), a centrally-located nucleotide-binding and oligomerization domain (NACHT) and C-terminal leucine-rich repeats (LRR). This gene product has a demonstrated role as a negative regulator of autophagy and type I interferon signaling pathways as a result of protein interactions with its NACHT domain. The PYD domain has also been shown to be important in the inhibition of NF-kB

(nuclear factor kappa-light-chain-enhancer of activated B cells). [provided by RefSeq, Dec 2016]

Applications: WB (1:500-2000)


Reactivity: Human

113 kDa MW.:

Subcellular Extracellular matrix

Location: ,Cytoplasm

VALIDATION IMAGES -



Sample: Lane 1: Human U-2 OS cell lysates Primary: Anti-NALP4 (bs-25219R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 113 kDa Observed band size: 120 kDa