

bs-12188R**[Primary Antibody]****ST18 Rabbit pAb**


Bioss
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GenelD:** 9705**Target:** ST18**Immunogen:** KLH conjugated synthetic peptide derived from human ST18/Znf387: 451-550/1047.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** ST18 is a 1,047 amino acid nuclear protein that is expressed at low levels in heart, liver, kidney, skeletal muscle, pancreas, testis, ovary and prostate. Containing six C2HC-type zinc fingers and an SMC domain, ST18 is a repressor protein that binds to DNA sequences containing a specific bipartite element. ST18 is considered a transcription factor that represses basal transcription activity from target promoters and inhibits colony formation in cultured breast cancer cells. ST18 is encoded by a gene located on human chromosome 8, which consists of nearly 146 million base pairs, encodes over 800 genes and is associated with a variety of diseases and malignancies. Schizophrenia, bipolar disorder, Trisomy 8, Pfeiffer syndrome, congenital hypothyroidism, Waardenburg syndrome and some leukemias and lymphomas are thought to occur as a result of defects in specific genes that map to chromosome 8.**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (3ug/Test)**Reactivity:** Human, Mouse, Rat
(predicted: Sheep, Dog, Horse)**Predicted MW:** 115 kDa**Subcellular Location:** Nucleus**— VALIDATION IMAGES —**


Sample: Cerebrum (Mouse) Lysate at 40 ug
Cerebrum (Rat) Lysate at 40 ug
Primary: Anti-ST18 (bs-12188R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 115 kD
Observed band size: 110 kD



Sample: U87MG(Human) Cell Lysate at 30 ug
Primary: Anti-ST18 (bs-12188R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 115 kD
Observed band size: 115 kD



Tissue/cell: rat brain tissue; 4%
Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-ST18 Polyclonal Antibody, Unconjugated(bs-12188R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: Mouse kidney. Primary Antibody

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

(green line): Rabbit Anti-ST18 antibody
(bs-12188R) Dilution: 3µg /10⁶ cells; Isotype
Control Antibody (orange line): Rabbit IgG .
Secondary Antibody : Goat anti-rabbit IgG-PE
Dilution: 1µg /test. Protocol The cells were
incubated in 5%BSA to block non-specific
protein-protein interactions for 30 min at at
room temperature .Cells stained with Primary
Antibody for 30 min at room temperature. The
secondary antibody used for 40 min at room
temperature. Acquisition of 20,000 events was
performed.