

bs-12204R**[Primary Antibody]****SALL4 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 57167 Target: SALL4 Immunogen: KLH conjugated synthetic peptide derived from human SALL4: 591-690/1053. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Sall4 is mammalian homologs of the Drosophila region-specific homeotic gene spalt, which encodes a zinc finger-containing transcription regulator. Drosophila spalt is an essential genetic component required for the specification of posterior head and anterior tail as opposed to trunk. Sall3 is expressed at 24 weeks of gestation in several regions of the human fetal brain including neurons of the hippocampus formation and of mediodorsal and ventrolateral thalamic nuclei, Purkinje cells of the cerebellum, and a subset of neurons in the brainstem. Sall4 expression in early mouse embryos is gradually confined to the head region and the primitive streak, followed by prominent expression in the developing midbrain, branchial arches, limbs and genital papilla.	Isotype: IgG SWISS: Q9UJQ4	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Pig, Sheep, Cow, Chicken, Dog, Horse) Predicted MW.: 112 kDa Subcellular Location: Nucleus
--	---	--

— VALIDATION IMAGES —

Sample: Lane 1: Mouse Testis tissue lysates Lane 2: Rat Testis tissue lysates Lane 3: Human Jurkat cell lysates Primary: Anti-SALL4 (bs-12204R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 112 kDa Observed band size: 118 kDa



Tissue/cell: rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-SALL4 Polyclonal Antibody, Unconjugated(bs-12204R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=2.775]** Ru Yu. et al. ATF6 deficiency damages the development of spermatogenesis in male Atf6 knockout mice. 2021 Dec 13 IHC ;Mouse. 34904262
- **[IF=2.4]** Zhao Xuan. et al. CircSfmbt2 Plays an Oncogenic Role in Lung Adenocarcinoma Depending on the miR-1305/SALL4 Axis. BIOCHEM GENET. 2023 Dec;;1-19 WB ;Human. 38127171

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.