### bsm-60019R

## [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# Arginase 1 Recombinant Rabbit mAb

DATASHEET -

Host: Rabbit Isotype: IgG Clonality: Recombinant CloneNo.: A6F1 **GenelD: 383 SWISS:** P05089

Target: Arginase 1

Immunogen: A synthesized peptide derived from human Arginase 1: 280-322.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Arginase I which is expressed almost exclusively in the liver, catalyzes the conversion of arginine to ornithine and urea . The human arginase I gene, which maps to chromosome 6q23, encodes a 322 amino acid protein. Arginase I exists as a homotrimeric protein and contains a binuclear manganese cluster. Arginase II catalyzes the same reaction as arginase I, but differs in its tissue specificity and subcellular location. Specifically, arginase II localizes to the mitochondria. Arginase II is expressed in nonhepatic tissues, with the highest levels of expression in the kidneys, but, unlike arginase I, is not expressed in liver. The human arginase II gene, which maps to chromosome 14q24.1-q24.3, encodes a 354 amino acid protein. In addition, arginase II contains a putative amino-terminal mitochondrial localization sequence.

Applications: WB (1:500-2000)


**IHC-P** (1:200-1000) **IHC-F** (1:400-800) **IF** (1:100-500)

Reactivity: Human, Mouse


Predicted 35 kDa

Subcellular Cytoplasm Location:


## VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with Arginase 1 monoclonal antibody, unconjugated (bsm-60019R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Arginase 1 Monoclonal Antibody, Unconjugated(bsm-60019R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Arginase 1 Monoclonal Antibody,

Unconjugated(bsm-60019R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.