bs-0957R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

PAG608 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 64393 SWISS: Q9HA38

Target: PAG608

Immunogen: KLH conjugated synthetic peptide derived from human WIG-1:

51-150/289.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a protein containing three zinc finger domains and a nuclear localization signal. The mRNA and the protein of this gene are upregulated by wildtype p53 and overexpression of this gene inhibits tumor cell growth, suggesting that this gene may have a role in the p53-dependent growth regulatory pathway. Alternative splicing of this gene results in two transcript variants encoding two isoforms differing in only one amino acid.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Human, Rat


(predicted: Mouse, Cow, Chicken, Dog, GuineaPig,

Horse)


Predicted 32 kDa MW.:

Subcellular Nucleus


VALIDATION IMAGES



Sample: Lane 1: 293T (Human) Cell Lysate at 30 ug Lane 2: Huvec (Human) Cell Lysate at 30 ug Lane 3: U87MG (Human) Cell Lysate at 30 ug Lane 4: U2os (Human) Cell Lysate at 30 ug Primary: Anti- PAG608 (bs-0957R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 35 kD Observed band size: 32 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PAG608) Polyclonal Antibody, Unconjugated (bs-0957R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PAG608) Polyclonal Antibody, Unconjugated (bs-0957R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.