bs-9870R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Thymosin beta 4 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 7114 SWISS: P62328

Target: Thymosin beta 4

Immunogen: KLH conjugated synthetic peptide derived from human Thymosin

beta 4: 37-44/44.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Proteins in the thymosin beta family are highly conserved polar peptides that bind monomeric Actin and thereby inhibit Actin polymerization. These proteins act as the main intracellular G-Actin sequestering peptides. The most abundant thymosin beta family member in mammalian cells and tissues is thymosin beta-4 (Tbeta-4), also designated Seraspenide. T beta-4 participates in several cellular events including cancerogenesis, apoptosis, angiogenesis, blood coagulation and would healing. Specifically, T beta-4 promotes cell migration and adhesion, accelerates wound healing and reduces inflammation, and becomes upregulated in a wide variety of human carcinomas. Due to the effects of T beta-4 in these events, it may become a protein of significant biological and pharmaceutical relevance.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:50-200)

Reactivity: Mouse (predicted: Human,

Rat)

Predicted 4.8 kDa

Subcellular Cytoplasm

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Thymosin beta 4) Polyclonal Antibody, Unconjugated (bs-9870R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.