

bs-0369M

[ Secondary Antibodies ]



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

## Mouse Anti-Rabbit IgM

### — DATASHEET —

<p><b>Host:</b> Mouse</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>Target:</b> Mouse Anti-Rabbit IgM</p> <p><b>Immunogen:</b> Native Rabbit IgM: full length.</p> <p><b>Purification:</b> affinity purified by Protein G</p> <p><b>Storage:</b> 0.01M PBS (pH7.4). Store at -20°C stable for 2 years (lyophilized powder). Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> Immunoglobulin M (IgM) normally constitutes about 10% of serum immunoglobulins. IgM antibody is prominent in early immune responses to most antigens and is largely confined to plasma due to its large size. Monomeric IgM is expressed as a membrane bound antibody on the surface of B cells and as a pentamer when secreted by plasma cells. Due to its high valency IgM is more efficient than other isotypes in binding antigens with repeating epitopes (virus particles and red blood cells) and is more efficient than IgG in activating the complement pathway. The gene for the mu constant region contains four domains separated by short intervening sequences.</p>	<p><b>Isotype:</b> IgG</p> <p><b>Applications:</b> Isotype Control Blocking Assay etc. Conjugate-Dependent.</p> <p><b>Reactivity:</b> Rabbit</p>
--	--

### — SELECTED CITATIONS —

- **[IF=4.8]** Wenpu Ren. et al. Salidroside treatment decreases the susceptibility of atrial fibrillation in diabetic mice by reducing mTOR-STAT3-MCP-1 signaling and atrial inflammation. INT IMMUNOPHARMACOL. 2024 Dec;142:113196 IF ;Mouse. 39306893