### bs-1368R

## [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

# SSTR3 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GeneID:** 6753 **SWISS:** P32745

Target: SSTR3

Immunogen: KLH conjugated synthetic peptide derived from human SSTR3:

121-230/418.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a member of the somatostatin receptor protein family. Somatostatins are peptide hormones that regulate diverse cellular functions such as neurotransmission, cell proliferation, and endocrine signaling as well as inhibiting the release of many hormones and other secretory proteins. Somatostatin has two active forms of 14 and 28 amino acids. The biological effects of somatostatins are mediated by a family of G-protein coupled somatostatin receptors that are expressed in a tissue-specific manner. Somatostatin receptors form homodimers and heterodimers with other members of the superfamily as well as with other G-protein coupled receptors and receptor tyrosine kinases. This protein is functionally coupled to adenylyl cyclase. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]

Applications: WB (1:500-2000)

400-901-9800

**IHC-P** (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Human, Mouse, Rat


(predicted: Cow, Dog,

Horse)


Predicted MW.: 46 kDa

Subcellular Location: Cell membrane


### VALIDATION IMAGES



Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Mouse Cerebellum tissue lysates Lane 3: Rat Cerebrum tissue lysates Lane 4: Human U87MG cell lysates Primary: Anti-SSTR3 (bs-1368R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 46 kDa Observed band size: 46 kDa



Tissue/cell: Mouse pancreas tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-SSTR3 Polyclonal Antibody, Unconjugated(bs-1368R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-SSTR3 Polyclonal Antibody, Unconjugated(bs-1368R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining