

**bs-12705R****[ Primary Antibody ]****MRPS22 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

|  |                      |  |
|--|----------------------|--|
| <b>Host:</b> Rabbit  | <b>Isotype:</b> IgG  | <b>Applications:</b> <b>WB</b> (1:500-2000)<br><b>IHC-P</b> (1:100-500)<br><b>IHC-F</b> (1:100-500)<br><b>IF</b> (1:100-500)<br><b>ICC/IF</b> (1:100-500)<br><b>ELISA</b> (1:5000-10000)<br><br><b>Reactivity:</b> (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)<br><br><b>Predicted MW.:</b> 41 kDa<br><br><b>Subcellular Location:</b> Cytoplasm |
| <b>Clonality:</b> Polyclonal   |                      |  |
| <b>GeneID:</b> 56945   | <b>SWISS:</b> P82650 |  |
| <b>Target:</b> MRPS22  |                      |  |
| <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human MRPS22: 151-250/360.   |                      |  |
| <b>Purification:</b> affinity purified by Protein A  |                      |  |
| <b>Concentration:</b> 1mg/ml   |                      |  |
| <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.<br>Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.   |                      |  |
| <b>Background:</b> Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that does not seem to have a counterpart in prokaryotic and fungal-mitochondrial ribosomes. This gene lies telomeric of and is transcribed in the opposite direction from the forkhead box L2 gene. A pseudogene corresponding to this gene is found on chromosome Xq. [provided by RefSeq] |                      |  |