bs-7562R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-TIRAP (Tyr86) Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 114609 **SWISS:** P58753

Target: TIRAP (Tyr86)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

TIRAP around the acetylation site of Tyr86: KD(p-Y)DV.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The Toll-like receptor (TLR) family in mammals comprises a family of transmembrane proteins characterized by multiple copies of leucine rich repeats in the extracellular domain and IL-1 receptor motif in the cytoplasmic domain. Like its counterparts in Drosophila, TLRs signal through adaptor molecules. The TLR family is a phylogenetically conserved mediator of innate immunity that is essential for microbial recognition. Ten human homologs of TLRs (TLR1-10) have been described. TIRAP (TIR domaincontaining adaptor protein) is an adaptor protein used by TLR4. Blocking TIRAP inhibits TLR4-mediated signaling events, including DC maturation and cytokine production.

Applications: WB (1:500-2000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) Flow-Cyt (2ug/Test)

Reactivity: Human, Rat


(predicted: Mouse, Pig, Cow, Dog, GuineaPig,

Horse)

Predicted 24 kDa MW.:

Subcellular Location: Cytoplasm

VALIDATION IMAGES




Sample: Lane 1: Human HepG2 cell lysates Lane 2: Human HUVEC cell lysates Lane 3: Human U251 cell lysates Lane 4: Human SH-SY5Y cell lysates Lane 5: Human K562 cell lysates Primary: Anti-phospho-TIRAP (Tyr86) (bs-7562R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 24 kDa Observed band size: 35 kDa



Paraformaldehyde-fixed, paraffin embedded (human heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (phospho-TIRAP (Tyr86)) Polyclonal Antibody, Unconjugated (bs-7562R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (phospho-TIRAP (Tyr86)) Polyclonal Antibody, Unconjugated (bs-7562R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Blank control: U937, Primary Antibody (green line): Rabbit Anti-phospho-TIRAP(Tyr86) antibody (bs-7562R) Dilution: 2µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-PE Dilution: $1\mu g$ /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block nonspecific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

• [IF=4.52] Rolf, Nina, et al. "Heterodimer - specific TLR2 stimulation results in divergent functional outcomes in B - cell precursor acute lymphoblastic leukemia." European journal of immunology (2015). Other; Human. 25867213