

Wnt3a Rabbit pAb

Catalog Number: bs-1700R

Target Protein: Wnt3a

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Rat (predicted:Human, Mouse, Rabbit, Pig, Sheep, Cow, Dog)

Predicted MW: 37 kDa

Entrez Gene: 89780

Swiss Prot: P56704

Source: KLH conjugated synthetic peptide derived from human Wnt3a: 281-352/352.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It encodes a protein which shows 96% amino acid identity to mouse Wnt3A protein, and 84% to human WNT3 protein, another WNT gene product. This gene is clustered with WNT14 gene, another family member, in chromosome 1q42 region. [provided by RefSeq, Jul 2008]

VALIDATION IMAGES



Sample: Lane 1: Cerebrum (Rat) Lysate at 40 ug Lane 2: Stomach (Rat) Lysate at 40 ug Primary: Anti-Wnt3a (bs-1700R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 39 kD Observed band size: 39 kD

PRODUCT SPECIFIC PUBLICATIONS

[IF=13.273] Xinkun Shen. et al. Improvement of aqueous stability and anti-osteoporosis properties of Zn-MOF coatings on titanium implants by hydrophobic raloxifene. Chem Eng J. 2021 Oct;:133094 WB ; Mouse . 10.1016/j.cej.2021.133094

[IF=13.12] Böske, Robert, et al. "Wnt Signaling in Chronic Rhinosinusitis with Nasal Polyps." American Journal of Respiratory Cell and Molecular Biology ja (2017). ICC ; ="Human" . 28059551

[IF=5.97] Sun et al. miR-491-5p, mediated by Foxi1, functions as a tumor suppressor by targeting Wnt3a/ β -catenin signaling in the development of gastric cancer. (2017) Cell.Death.Dis. 8:e2714 IHC,WB ; Human . 28358374

[IF=4.6] Ippei Horibe. et al. Acquired curved hair is caused by fusion of multiple hair matrix cells. J DERMATOL SCI. 2024 Feb;: IHC ; Human . 38431439

[IF=3.8] Mingyi Huo. et al. Huaier promotes sensitivity of colorectal cancer to oxaliplatin by inhibiting METTL3 to regulate the Wnt/ β -catenin signaling pathway. ONCOL REP. 2025 Jan;53(1):1-13 WB ; Human . 39513580