

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## DPP10 Rabbit pAb

Catalog Number: bs-13028R

Target Protein: DPP10
Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:100-500),

ELISA (1:5000-10000)

Reactivity: (predicted:Human, Mouse, Rat, Horse)

Predicted MW: 91 kDa Entrez Gene: 57628 Swiss Prot: Q8N608

Source: KLH conjugated synthetic peptide derived from human DPP10: 631-730/796.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Dipeptidyl peptidases (DPPs) mediate regulatory activity of their substrates and have been

linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs can bind specific voltage-gated potassium channels and alter their

expression and biophysical properties and may also influence T cells. DPP proteins include DPRP1, DPRP2, DPP3, DPP7, DPP10, DPPX and CD26. DPP10 (dipeptidyl-peptidase 10), also

known as DPRP3 (dipeptidyl peptidase IV-related protein 3), DPL2 or DPPY, is a non-

functional dipeptidyl peptidase which can bind to the potassium channels KV4.1 and KV4.2.

It is a single-pass type II membrane protein expressed in spinal cord, adrenal glands, pancreas and brain tissues and may act as a modulator for cell surface expression and

activity of KV4.1 and KV4.2.