

bs-13028R**[Primary Antibody]****DPP10 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 57628**SWISS:** Q8N608**Target:** DPP10**Immunogen:** KLH conjugated synthetic peptide derived from human DPP10: 631-730/796. < Extracellular >**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Dipeptidyl peptidases (DPPs) mediate regulatory activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs can bind specific voltage-gated potassium channels and alter their expression and biophysical properties and may also influence T cells. DPP proteins include DPRP1, DPRP2, DPP3, DPP7, DPP10, DPPX and CD26. DPP10 (dipeptidyl-peptidase 10), also known as DPRP3 (dipeptidyl peptidase IV-related protein 3), DPL2 or DPPY, is a non-functional dipeptidyl peptidase which can bind to the potassium channels KV4.1 and KV4.2. It is a single-pass type II membrane protein expressed in spinal cord, adrenal glands, pancreas and brain tissues and may act as a modulator for cell surface expression and activity of KV4.1 and KV4.2.

Applications: WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**ICC/IF** (1:100-500)**ELISA** (1:5000-10000)**Reactivity:** (predicted: Human, Mouse, Rat, Horse)**Predicted MW.:** 91 kDa**Subcellular Location:** Cell membrane