

**bs-19698R****[ Primary Antibody ]****QSOX2 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)  <b>Reactivity:</b> Mouse (predicted: Human, Rat)  <b>Predicted MW.:</b> 76 kDa  <b>Subcellular Location:</b> Secreted ,Cell membrane ,Nucleus
<b>Clonality:</b> Polyclonal		
<b>GenelD:</b> 169714	<b>SWISS:</b> Q6ZRP7	
<b>Target:</b> QSOX2		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human QSOX2: 101-200/698.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> QSOX2 is a member of the sulfhydryl oxidase/quiescin-6 (Q6) family (QSOX1; MIM 603120) that regulates the sensitization of neuroblastoma cells for IFN-gamma (IFNG; MIM 147570)-induced cell death (Wittke et al., 2003 [PubMed 14633699]).[supplied by OMIM, Jun 2009]		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (QSOX2) Polyclonal Antibody, Unconjugated (bs-19698R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.