
MMP14 Antibody Blocking Peptide

Catalog Number: bs-0414P

Activity: Not tested

Purification: HPLC

Storage: Shipped at 4°C. Stored at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: MMP14 may be an activator of pro gelatinase A and is expressed in fibroblast cells during both wound healing and human cancer progression. MMP14 is expressed in very low levels and may require stimulation with concanavolin A or the phorbol ester TPA to stimulate production of MMP14.

Matrix metalloproteinase-14 precursor is endopeptidase that degrades various components of the extracellular matrix, such as collagen. Activates progelatinase A. Essential for pericellular collagenolysis and modeling of skeletal and extraskelatal connective tissues during development. [Catalytic activity] Endopeptidase activity. Activates progelatinase A by cleavage of the propeptide at 37-Asn-|-Leu-38. Other bonds hydrolyzed include 35-Gly-|-Ile-36 in the propeptide of collagenase 3, and 341-Asn-|-Phe-342, 441-Asp-|-Leu-442 and 354-Gln-|-Thr-355 in the aggrecan interglobular domain. Highly expressed in placenta, kidney, heart, lung, embryonic skeletal and periskeletal tissues. Belongs to the peptidase M10A family.