

bs-5699R**[Primary Antibody]****phospho-FADD (Ser194) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 8772	SWISS: Q13158	IF (1:100-500)
Target: phospho-FADD (Ser194)		Reactivity: Human
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human FADD around the phosphorylation site of Ser194: AM(p-S)PM.		
Purification: affinity purified by Protein A		Predicted MW.: 23 kDa
Concentration: 1mg/ml		Subcellular Location: Cytoplasm
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Predicted to enable several functions, including caspase binding activity; death effector domain binding activity; and tumor necrosis factor receptor superfamily binding activity. Involved in several processes, including hematopoietic or lymphoid organ development; negative regulation of activation-induced cell death of T cells; and positive regulation of CD8-positive, alpha-beta cytotoxic T cell extravasation. Acts upstream of or within extrinsic apoptotic signaling pathway in absence of ligand; motor neuron apoptotic process; and regulation of programmed cell death. Predicted to be located in several cellular components, including cell body; cytosol; and membrane raft. Predicted to be part of CD95 death-inducing signaling complex and ripoptosome. Predicted to be active in cytoplasm. Is expressed in several structures, including alimentary system; brain; genitourinary system; hemolymphoid system gland; and liver and biliary system. Human ortholog(s) of this gene implicated in leukemia. Orthologous to human FADD (Fas associated via death domain). [provided by Alliance of Genome Resources, Apr 2022]		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-FADD (Ser194)) Polyclonal Antibody, Unconjugated (bs-5699R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023)



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-FADD (Ser194)) Polyclonal Antibody, Unconjugated (bs-5699R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023)

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

instructionsand DAB staining.

instructionsand DAB staining.