

bs-9471R**[Primary Antibody]****MESP1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 55897	SWISS: Q9BRJ9	IHC-F (1:100-500)
Target: MESP1		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human MESP1: 101-200/268.		Reactivity: Human, Mouse, Rat (predicted: Pig, Cow, Chicken, Dog)
Purification: affinity purified by Protein A		Predicted MW.: 30 kDa
Concentration: 1mg/ml		Subcellular Location: Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: MESP1, also known as bHLHc5, is a 268 amino acid protein that contains one basic helix-loop-helix (bHLH) domain, a motif that mediates protein dimerization and can bind to the E-box sequence of DNA. Localized to the nucleus, MESP1 functions as a transcription factor that, via its bHLH domain, participates in the epithelialization and the development of the cardiac and somitic mesoderm. MESP1 is highly expressed during gastrulation and somitogenesis and is necessary for the formation of single heart tubes during cardiac maturation. Early detection of MESP1 may be an indicator of the formation of cardiac precursor cells in developing embryos. Additionally, MESP1 plays a role in the rostrocaudal patterning of the somites, an event that influences select Notch signaling pathways.		

— VALIDATION IMAGES —

Sample: Testis (Mouse) Lysate at 40 ug
Primary: Anti-MESP1 (bs-9471R) at 1/500 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 30 kD
Observed band size: 30 kD



Tissue/cell: Mouse embryos tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-MESP1 Polyclonal Antibody, Unconjugated(bs-9471R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: mouse stomach tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-MESP1 Polyclonal Antibody, Unconjugated(bs-9471R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining