

---

## Wnt16 Rabbit pAb

Catalog Number: bs-9476R

Target Protein: Wnt16

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human (predicted:Mouse, Rat, Rabbit, Pig, Cow)

Predicted MW: 37 kDa

Entrez Gene: 51384

Swiss Prot: Q9UBV4

Source: KLH conjugated synthetic peptide derived from human Wnt16: 186-365/365.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** The Wnt genes encode a family of secreted extracellular signaling glycoproteins, which function in a variety of important developmental processes such as regulation of cell growth and differentiation. Wnt proteins also play roles in carcinogenesis. Wnt-14, rather than Wnt-15, is preferentially expressed in various types of human cancer. Wnt-15 is expressed in fetal and adult kidney and is most homologous to Wnt-14. Wnt-16, another member in the Wnt family, has two mRNA isoforms, Wnt-16a and Wnt-16b. These isoforms differ in the composition of their 5'UTR and first exon, which results in differential expression. Wnt-16a is expressed only in pancreas, whereas Wnt-16b is highly expressed in adult kidney, placenta, brain, heart and spleen, but not in bone marrow. However, Wnt-16 transcripts are present in bone marrow and cell lines derived from pre-B acute lymphoblastoid leukemia patients carrying the E2A-Pbx1 hybrid gene. Thus, Wnt-16 is a downstream target of E2A-Pbx1, and the Wnt-16-mediated autocrine growth mechanism may contribute to the development of t(1;19) pre-B acute lymphoblastoid leukemias.

### VALIDATION IMAGES

---



Sample: A431(Human) Cell Lysate at 30 ug Primary: Anti- Wnt16 (bs-9476R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 37 kD Observed band size: 56 kD