bs-23474R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Nano-Tag (15) Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: Nano-Tag (15)

Purification: affinity purified by Protein A

Concentration: 1mg/1ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Well-characterized antibodies for epitope tags consisting of short

sequences are widely used in the study of protein expression in various systems. The Nano-tag is a new streptavidin-binding peptide for both the purification and the detection of Nano-tagged


proteins. This peptide possesses nanomolar-affinity for strents vidin and therefore is termed Nano-tag. The nano-tags

streptavidin and therefore is termed Nano-tag. The nano-tags have two types, Nano-tag15 (MDVEAWLGARVPLVET) and Nano-tag9 (MDVEAWLGAR), which bind to streptavidin with dissociation

constants of 4 nM and 17 nM, respectively.

constants of

VALIDATION IMAGES



Sample: Lane 1: Nano-Tagged Fusion Protein Overexpression E.coli Lysate (Cat#: bs-41403P) at 2ug Lane 2: Nano-Tagged Fusion Protein Overexpression E.coli Lysate (Cat#: bs-41403P) at 4ug Primary: Anti-Nano-Tag15 (bs-23474R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51 kD Observed band size: 51 kD **Applications: WB** (1:1000-2000)

IHC-P (1:200-1000)
IHC-F (1:200-1000)
IF (1:200-1000)
Flow-Cyt (1ug/Test)
ICC/IF (1:100-500)
ELISA (1:1000-5000)

Reactivity: Species independent