

bs-13597R**[Primary Antibody]****HESX1 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 8820 Target: HESX1 Immunogen: KLH conjugated synthetic peptide derived from human HESX1: 51-150/185. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The homeobox protein, HESX1, which is also known as Rathke's pouch homeobox, HANF, homeodomain transcription factor, and anterior-restricted homeobox protein is a transcription factor that belongs to the homeodomain family of DNA binding proteins. HESX1 is initially expressed in embryonic stem cells and the primitive forebrain, and is essential for normal development of the eyes and other anterior CNS structures, such as the hypothalamus, the pituitary gland and the olfactory bulbs. The homeobox gene Hesx1 is expressed in the anterior visceral endoderm (AVE), anterior axial mesendoderm (AME), and anterior neural ectoderm (ANE) during early embryogenesis. Mutations in the Hesx1 gene are associated with disorders that are comparable with septo-optic dysplasia (SOD). These disorders are characterized by hypoplasia of the optic nerve, various types of forebrain defects and pituitary hormone deficiencies, including hypothyroidism. Hesx1 also acts as a transcriptional repressor of reporter gene constructs in tissue culture assays.	Isotype: IgG SWISS: Q9UBX0 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Rat (predicted: Human, Mouse, Rabbit, Pig, Sheep, Cow, Dog, Horse) Predicted MW.: 21 kDa Subcellular Location: Nucleus
--	---

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HESX1) Polyclonal Antibody, Unconjugated (bs-13597R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.