## bs-13606R

## [ Primary Antibody ]

## BCL6B Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500)
Clonality: Polyclonal		<b>IF</b> (1:100-500)
GenelD: 255877	SWISS: Q8N143	ICC/IF (1:100-500)
Target: BCL6B		ELISA (1:5000-10000)
Immunogen: KLH conjugated synthetic peptide derived from human BCL6B: 141-440/479.		<b>Reactivity:</b> (predicted: Human, Mouse, Rat, Pig, Sheep, Cow, Dog)
Purification: affinity purified by I	Protein A	
Concentration: 1mg/ml		Prodicted
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated		Predicted MW.: <sup>51 kDa</sup> Subcellular
freeze/thaw cycles.		Subcellular Location: Nucleus
elements and influe differentiation. Add expression, thereby Bcl-6b (B-cell CLL/I BAZF or ZBTB28, is one BTB (POZ) dom ubiquitously with h associates with Bcl transcriptional repr B-cell development	nal repressor, binds Stat recognition-like DNA ences germinal center development and cell itionally, Bcl-6 negatively regulates NF x B inhibiting NF x B-mediated cellular functions. /mphoma 6, member B), also known as ZNF62, a 480 amino acid nuclear protein that contains ain and five C2H2-type zinc fingers. Expressed ighest expression in placenta and heart, Bcl-6b 6 and functions as a sequence-specific essor that is thought to be necessary for early . The gene encoding Bcl-6b maps to human nich comprises over 2.5% of the human genome ,200 genes.	

## - SELECTED CITATIONS -

• [IF=1.813] Zhang Yu. et al. The Combination of Jiedu Xiaoluo Decoction with Autologous Peripheral Blood Stem Cell Transplantation (APBSCT) Accelerates Disease Remission of Non-Hodgkin Lymphoma. Evid-Based Compl Alt.

2021;2021:2745705 IHC ;Mouse. 33505491