

bsm-60738R**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Ki67 Recombinant Rabbit mAb


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:200-800) IHC-F (1:200-800) IF (1:200-800) Flow-Cyt (1:50-100) ICC/IF (1:50-200)
Clonality: Recombinant	CloneNo.: 6B9	
GeneID: 4288	SWISS: P46013	
Target: Ki67		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Reactivity: Human, Mouse, Rat
Storage: PBS, Glycerol, BSA. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Ki67 antigen is the prototypic cell cycle related nuclear protein, expressed by proliferating cells in all phases of the active cell cycle (G1, S, G2 and M phase). It is absent in resting (G0) cells. Ki67 antibodies are useful in establishing the cell growing fraction in neoplasms (immunohistochemically quantified by determining the number of Ki67 positive cells among the total number of resting cells = Ki67 index). In neoplastic tissues the prognostic value is comparable to the tritiated thymidine labelling index. The correlation between low Ki67 index and histologically low grade tumours is strong. Ki67 is routinely used as a neuronal marker of cell cycling and proliferation.		
		Predicted MW.: 358 kDa Subcellular Location: Nucleus


— VALIDATION IMAGES —




Paraformaldehyde-fixed, paraffin embedded (mouse thymus); Antigen retrieval by boiling in sodium EDTA buffer (Ph9.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Ki67) Monoclonal Antibody, Unconjugated (bsm-60738R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (rat spleen); Antigen retrieval by boiling in sodium EDTA buffer (Ph9.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Ki67) Monoclonal Antibody, Unconjugated (bsm-60738R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




4% Paraformaldehyde-fixed HeLa(H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (Ki-67) monoclonal Antibody, unconjugated (bsm-60738R) 1:200, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.



Paraformaldehyde-fixed, paraffin embedded Human Esophageal Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Ki67 Monoclonal Antibody, Unconjugated (bsm-60738R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-BF488), DAPI (blue, C02-04002) was used to stain



Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Ki67 Monoclonal Antibody, Unconjugated (bsm-60738R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-BF488), DAPI (blue, C02-04002) was used to stain



Blank control (black line) :HeLa. Primary Antibody (green line): Rabbit Anti-Ki67 antibody (bsm-60738R) Dilution:0.5ug/Test; Secondary Antibody (white/blue line) : Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90%

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

the cell nuclei.

the cell nuclei.

ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=13.3]** Jie Meng. et al. Hollow and heterojunction-structured Janus nanomotors for pyroelectrodynamic and nanozyme-catalyzed tumor therapy. CHEM ENG J. 2024 Dec;501:157763 IHC ;Mouse. 10.1016/j.cej.2024.157763
- **[IF=10.6]** Tian Yafei. et al. Biomimetic mesenchymal stem cell membrane-coated nanoparticle delivery of MKP5 inhibits hepatic fibrosis through the IRE/XBP1 pathway. J NANOBIOTECHNOL. 2024 Dec;22(1):1-25 IF ;Mouse. 39609656
- **[IF=10.5]** Chen Yang. et al. A pH and glutathione-responsive carbon monoxide-driven nano-herb delivery system for enhanced immunotherapy in colorectal cancer. J CONTROL RELEASE. 2024 Dec;376:659 IHC ;Mouse. 39442888
- **[IF=10.8]** Kaili Deng. et al. Chemo-photothermal nanoplatform with diselenide as the key for ferroptosis in colorectal cancer. J CONTROL RELEASE. 2024 Feb;366:684 IHC ;Mouse. 38224739
- **[IF=10.6]** Li Kangkang. et al. Reactive oxygen species/glutathione dual sensitive nanoparticles with encapsulation of miR155 and curcumin for synergized cancer immunotherapy. J NANOBIOTECHNOL. 2024 Dec;22(1):1-23 IHC ;Mouse. 38972995