


bsm-52469R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

CD4 Recombinant Rabbit mAb**— DATASHEET —****Host:** Rabbit**Clonality:** Recombinant**GeneID:** 920**Target:** CD4**Immunogen:** A synthesized peptide derived from human CD4: 200-400.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** This gene encodes a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigens and is also a receptor for the human immunodeficiency virus. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene. [provided by RefSeq, Aug 2010].**Isotype:** IgG**CloneNo.:** 65H6**SWISS:** P01730**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (1ug/Test)**Reactivity:** Human**Predicted**
MW.: 48 kDa**Subcellular**
Location: Cell membrane**— VALIDATION IMAGES —**


25 ug total protein per lane of various lysates (see on figure) probed with CD4 monoclonal antibody, unconjugated (bsm-52469R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Human Gastric Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD4 Monoclonal Antibody, Unconjugated (bsm-52469R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD4 Monoclonal Antibody, Unconjugated (bsm-52469R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD4 Monoclonal Antibody, Unconjugated (bsm-52469R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-02965-



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD4 Monoclonal Antibody, Unconjugated (bsm-52469R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-02965-



Blank control: Jurkat. Primary Antibody (green line): Rabbit Anti-CD4 antibody (bsm-52469R) Dilution: 1ug/Test; Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line): Normal Rabbit IgG Protocol The cells were incubated in 5%BSA to block non-specific

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei.

BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei.

protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=18.027]** Zhu Li. et al. Immunogenic Cell Death Augmented by Manganese Zinc Sulfide Nanoparticles for Metastatic Melanoma Immunotherapy. ACS NANO. 2022;XXXX(XXX):XXX-XXX IHC ;Mouse. 35981098
- **[IF=7.59]** Guoyun Wan. et al. Endoplasmic reticulum-targeted NIR-II phototherapy combined with inflammatory vascular suppression elicits a synergistic effect against TNBC. BIOMATER SCI-UK. 2023 Jan;; IF ;Mouse. 36692120
- **[IF=7.3]** Liquan Jin. et al. High expression ITGA2 affects the expression of MET, PD-L1, CD4 and CD8 with the immune microenvironment in pancreatic cancer patients. FRONT IMMUNOL. 2023; 14: 1209367 IHC ;Human. 37881431
- **[IF=7.1]** Shuangjie Zhu. et al. Dihydroartemisinin inhibits the development of autoimmune thyroiditis by modulating oxidative stress and immune imbalance. free radical biology and medicine. 2025 Feb 21:231:57-67. IF ;Mouse. 39988064
- **[IF=3.3]** Jiayun Jiang. et al. The immune-reinforcements of Lenvatinib plus anti-PD-1 and their rationale to unite with TACE for unresectable hepatocellular carcinoma treatment. IMMUNOL LETT. 2025 Oct;275:107003 IF ;Human,Mouse. 40189154