

bsm-52304R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Bcl-2 Recombinant Rabbit mAb**— DATASHEET —**

Host: Rabbit Clonality: Recombinant GeneID: 596 Target: Bcl-2 Immunogen: A synthesized peptide derived from human Bcl 2: 48-81. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The Bcl-2 gene was isolated at the chromosomal breakpoint of t(14;18)-bearing follicular B cell lymphomas(1,2).Bcl-2 blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal (3,5).Bcl-2 appears to function in several subcellular locations yet lacks any known motifs that would confer insight into its mechanism of action (6,7).A more recently identified protein,designated Bax p21(i.e., Bcl-associated X protein),has extensive amino acid homology with Bcl-2 and both homodimerizes and forms heterodimers with Bcl-2(8). Overexpression of Bax accelerates apoptotic death induced by cytokine deprivation in an IL-3 dependent cell line and Bax also counters the death repressor activity of Bcl-2(8).	Isotype: IgG CloneNo.: 5F7 SWISS: P10415	Applications: WB (1:500-2000) Reactivity: Human Predicted MW.: 26 kDa Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus
--	---	--

— VALIDATION IMAGES —

25 ug total protein per lane of various lysates (see on figure) probed with Bcl-2 monoclonal antibody, unconjugated (bsm-52304R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

— SELECTED CITATIONS —

- **[IF=7]** Zhang Zhichun. et al. A novel SLC25A1 inhibitor, parthenolide, suppresses the growth and stemness of liver cancer stem cells with metabolic vulnerability. CELL DEATH DISCOV. 2023 Sep;9(1):1-12 WB ;Human. 37741815
- **[IF=5.722]** Wang, Sanchun. et al. Pseudoginsengonin DQ exerts antitumour activity against hypopharyngeal cancer cells by targeting the HIF-1α-GLUT1 pathway. Cancer Cell Int. 2021 Dec;21(1):1-12 WB ;Human. 34281558
- **[IF=5.895]** Yu-Sheng Shi. et al. Pteris laeta Wall. and Its New Phytochemical, Pterosinsade A, Promote Hippocampal Neurogenesis via Activating the Wnt Signaling Pathway. J AGR FOOD CHEM. 2023;XXXX(XXX):XXX-XXX WB ;Murine. 36892329
- **[IF=5.3]** Caixia Tan. et al.Mcl-1 downregulation enhances BCG treatment efficacy in bladder cancer by promoting

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

macrophage polarization.Cancer Cell International.2025 Feb 15;25(1):48. ;. 39955585

- **[IF=4.5]** Gan Rao. et al. Arsenic and polystyrene-nano plastics co-exposure induced testicular toxicity: Triggers oxidative stress and promotes apoptosis and inflammation in mice. ENVIRON TOXICOL. 2023 Sep;; WB,IF ;Mouse. 37705229