

[Primary Antibody]

Trk A + B + C Recombinant Rabbit mAb



BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioass.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET

Host: Rabbit

Isotype: IgG

Clonality: Recombinant

CloneNo.: 5T6

GeneID: 4914

SWISS: P04629

Target: Trk A + B + C

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The Trk family of nerve growth factor receptors includes Trk A (also referred to as Trk A gp140), Trk B and Trk C. The prototype member of this gene family, Trk A, encodes a 140 kDa cell surface receptor, gp140, the expression of which is restricted in vivo to neurons of the sensory spinal and cranial ganglia of neurocrest origin. Nerve growth factor (NGF) stimulates tyrosine phosphorylation of Trk gp140 in neural cell lines and in embryonic dorsal root ganglia. By comparison, BDNF and to a lesser extent, NT-3, but not NGF, can induce tyrosine phosphorylation of Trk B gp145. The third member of the Trk receptor family, Trk C encodes a 140 kDa protein, Trk C gp140, that is preferentially expressed in brain tissue and primarily functions as a receptor for NT-3. An additional component of the Trk receptor complex, NGFR p175, binds to neurotrophic factors with low affinity but is required for efficient signaling. NGFR p175 accelerates Trk activation and may recruit downstream effector molecules to the ligand-bound receptor complex.

Applications: WB (1:500-2000)

IHC-P (1:50-200)

IHC-F (1:50-200)

IF (1:50-200)


ICC/IF (1:50-)

Reactivity: Human, Mouse, Rat


Predicted
MW.: 90 kDa

Subcellular Location: Cell membrane ,Cytoplasm


— VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with Trk A + B + C monoclonal antibody, unconjugated (bsm-52715R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Rat Cerebellum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Trk A + B + C Monoclonal Antibody, Unconjugated (bsm-52715R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Cerebellum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Trk A + B + C Monoclonal Antibody, Unconjugated (bsm-52715R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.