bs-4240R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn

techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

DDIT4 Rabbit pAb

GenelD: 54541 SWISS: Q9NX09

Target: DDIT4

Immunogen: KLH conjugated synthetic peptide derived from human DDIT4:

165-232/232.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Inhibits cell growth by regulating the TOR signaling pathway

upstream of the TSC1-TSC2 complex and downstream of AKT1.

Promotes neuronal cell death.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Rat (predicted: Human,


Mouse, Rabbit, Pig, Cow,

Horse)

Predicted MW.: 25 kDa

Subcellular Cytoplasm Location:

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat heart tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DDIT4) Polyclonal Antibody, Unconjugated (bs-4240R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

— SELECTED CITATIONS ————

• [IF=4.8] Yang Liu. et al. Aerobic exercise mitigates hippocampal neuronal apoptosis by regulating DAPK1/CDKN2A/REDD1/FoXO1/FasL signaling pathway in D-galactose-induced aging mice. FASEB J. 2023 Sep;37(10):e23205 WB; Mouse. 37768886