

Recombinant human Bcl-2 protein, C-His

Catalog Number: bs-10846P

Concentration: >0.5 mg/ml

AA Seq: 1-211/239

Predicted MW: 24

Tags: C-His

Activity: Not tested

Endotoxin: Not analyzed

Purity: >95% as determined by SDS-PAGE

Purification: AC

Form: Lyophilized or Liquid

Storage: 20mM Tris-HCl (pH7.5) with 8M Urea.

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: The Bcl-2 gene was isolated at the chromosomal breakpoint of t(14;18)-bearing follicular B cell lymphomas(1,2).Bcl-2 blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal (3,5).Bcl-2 appears to function in several subcellular locations yet lacks any known motifs that would confer insight into its mechanism of action (6,7).A more recently identified protein,designated Bax p21(i.e., Bcl-associated X protein),has extensive amino acid homology with Bcl-2 and both homodimerizes and forms heterodimers with Bcl-2(8). Overexpression of Bax accelerates apoptotic death induced by cytokine deprivation in an IL-3 dependent cell line and Bax also counters the death repressor activity of Bcl-2(8).

VALIDATION IMAGES



The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.