

bsm-54201R

[Primary Antibody]

HLA-DQA1 Recombinant Rabbit mAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

DATASHEET

Host: Rabbit

Clonality: Recombinant

Target: HLA-DQA1

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Isotype: IgG

CloneNo.: 8C1

Applications: WB (1:500-2000)

IHC-P (1:50-200)


IHC-F (1:50-200)

IF (1:50-200)


Reactivity: Human, Mouse, Rat

Predicted MW.: 26 kDa


VALIDATION IMAGES



Sample: Lane 1: Mouse Pancreas lysates Lane 2: Mouse Spleen lysates Lane 3: Mouse Lymph node lysates Lane 4: Rat Pancreas lysates Lane 5: Rat Spleen lysates Lane 6: Rat Lymph node lysates Lane 7: Human Raji Cell Lysates Primary: Anti- HLA-DQA1 (bsm-54201R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 26 kDa Observed band size: 30 kDa



Paraformaldehyde-fixed, paraffin embedded (mouse spleen); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HLA-DQA1) Monoclonal Antibody, Unconjugated (bsm-54201R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat spleen); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HLA-DQA1) Monoclonal Antibody, Unconjugated (bsm-54201R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.