
phospho-CDKN1B (Ser10) Rabbit pAb

Catalog Number: bs-20074R

Target Protein: phospho-CDKN1B (Ser10)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: **WB** (1:500-2000), **IHC-P** (1:100-500), **IHC-F** (1:100-500), **IF** (1:100-500), **Flow-Cyt** (1µg/Test)

Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Dog, GuineaPig)

Predicted MW: 22 kDa

Subcellular: Cytoplasm, Nucleus

Locations:

Entrez Gene: 1027

Swiss Prot: P46527

Source: KLH conjugated synthesised phosphopeptide derived from human CDKN1B around the phosphorylation site of Ser10: NG(p-S)PS.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Cell cycle progression is regulated by cyclins and their cognate Cdk. p27 KIP 1 is a cell cycle regulatory mitotic inhibitor of cdk activity. p27 KIP 1 is a candidate tumor suppressor gene, and has been proposed to function as a possible mediator of TGF beta induced G1 arrest. p27 KIP 1 is up regulated in response to antimitogenic stimuli. The increased protein expression of p27 results in cellular arrest by binding to cyclin/Cdk complexes such as cyclin D1/Cdk4. p27 Kip1 is regulated by phosphorylation on serine 10 (S10) and threonine 187 (T187). Phosphorylation by CDK2 on T187 results in ubiquitylation and degradation of p27 Kip 1; while phosphorylation by hKIS on S10 signals the nuclear export to the cytoplasm.


VALIDATION IMAGES




Sample: Lymphocyte (Rat) Lysate at 40 ug Primary: Anti- phospho-CDKN1B (Ser10) (bs-20074R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 22 kD
Observed band size: 22kD




Sample: MOLT-4 Cell (Human) Lysate at 30 ug Primary: Anti- phospho-CDKN1B (Ser10) (bs-20074R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 22 kD
Observed band size: 22kD




Sample: Muscle (Mouse) Lysate at 40 ug Primary: Anti- phospho-CDKN1B (Ser10) (bs-20074R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 22 kD
Observed band size: 22kD



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-CDKN1B (Ser10)) Polyclonal Antibody, Unconjugated (bs-20074R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-CDKN1B (Ser10)) Polyclonal Antibody, Unconjugated (bs-20074R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control (blue line): MCF7(fixed with 70% ethanol (Overnight at 4°C) and then permeabilized with 90% ice-cold methanol for 30 min on ice) Primary Antibody (green line): Rabbit Anti-phospho-CDKN1B (Ser10) antibody (bs-20074R), Dilution: 1µg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG .
Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC, Dilution: 1µg /test.