

bs-9447R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

BMP10 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: ELISA (1:5000-10000)
Clonality: Polyclonal		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow, Horse)
GeneID: 27302	SWISS: Q95393	
Target: BMP10		Predicted MW.: 12 kDa
Immunogen: KLH conjugated synthetic peptide derived from human BMP10: 331-424/424.		Subcellular Location: Secreted
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Bone morphogenic proteins (BMPs) are members of the TGF beta superfamily and are involved in the induction of cartilage and bone formation. BMP-10 (Bone morphogenetic protein 10) is a 424 amino acid secreted protein that exists as a disulfide-linked homodimer. Expressed specifically in heart tissue, BMP-10 functions to prevent the premature activation of p57 (a negative cell cycle regulator), thereby maintaining the proliferative activity of embryonic cardiomyocytes and playing a critical role in the formation of trabeculae within the developing heart. Additionally, BMP-10 inhibits the growth and migration of endothelial cells and functions as a ligand for ALK-1, a protein that is tightly linked to angiogenesis. This suggests that, in addition to its role in cardiomyocyte formation, BMP-10 participates in the formation of blood vessels in developing cardiac muscle.		

— SELECTED CITATIONS —

- **[IF=2.24]** Wu, Mei, et al. "Expression analysis of BMP2, BMP5, BMP10 in human colon tissues from Hirschsprung disease patients." Int J Clin Exp Pathol 7.2 (2014): 529-536. WB ;="Human". 24551273