

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

EEF1D Rabbit pAb

Catalog Number: bs-13055R

Target Protein: EEF1D Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse (predicted: Human, Rat, Rabbit, Pig, Sheep, Cow, Chicken, Horse)

Predicted MW: 31 kDa

Subcellular Cytoplasm, Nucleus

Locations:

Entrez Gene: 1936 Swiss Prot: P29692

Source: KLH conjugated synthetic peptide derived from human EEF1D: 1-100/281.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: This gene encodes a subunit of the elongation factor-1 complex, which is responsible for the

enzymatic delivery ofaminoacyl tRNAs to the ribosome. This subunit, delta, functions asguanine nucleotide exchange factor. It is reported that followingHIV-1 infection, this subunit interacts with HIV-1 Tat. This interaction results in repression of translation of host

cellproteins and enhanced translation of viral proteins. Severalalternatively spliced transcript variants encoding multipleisoforms have been found for this gene. Related pseudogenes have been defined on chromosomes 1, 6, 7, 9, 11, 13, 17, 19.[provided]

byRefSeq, Aug 2010].

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EEF1D) Polyclonal Antibody, Unconjugated (bs-13055R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.