

**bs-13062R****[ Primary Antibody ]****EGFL6 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ICC/IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)  <b>Reactivity:</b> (predicted: Human, Mouse, Rat, Sheep, Cow, Dog, Horse)  <b>Predicted MW.:</b> 59 kDa  <b>Subcellular Location:</b> Secreted ,Extracellular matrix ,Cell membrane
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 25975	<b>SWISS:</b> Q8IUX8	
<b>Target:</b> EGFL6		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human EGFL6: 231-330/553.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> This gene encodes a member of the epidermal growth factor(EGF) repeat superfamily. Members of this superfamily are characterized by the presence of EGF-like repeats and are often involved in the regulation of cell cycle, proliferation, and developmental processes. The gene product contains a signal peptide, suggesting that it is secreted; an EGF repeat region consisting of 4 complete EGF-like repeats and 1 partial EGF-like repeat, 3 of which have a calcium-binding consensus sequence; an arg-gly-asn integrin association motif; and a MAM domain, which is believed to have an adhesive function. This gene is expressed early during development, and its expression has been detected in lung and meningioma tumors. [provided by RefSeq, Jul 2008].		