

bs-9458R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

TNNI3K Rabbit pAb**— DATASHEET —**

Host: Rabbit Clonality: Polyclonal GeneID: 51086 Target: TNNI3K Immunogen: KLH conjugated synthetic peptide derived from human TNNI3K: 1-100/835. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: TNNI3K, also known as CARK, is a 936 amino acid serine/threonine-protein kinase that is highly expressed in heart. Overexpression of TNNI3K leads to improved cardiac function by enhancing beating frequency and increasing contractile force and epinephrine response. TNNI3K suppresses phosphorylation of cardiac troponin I and p38/JNK-mediated apoptosis, therefore protecting the myocardium from ischemic injury. Administration of TNNI3K to mice with myocardial infarction improves cardiac performance and attenuates ventricular remodeling, suggesting that TNNI3K could be a promising target in the treatment of cardiac diseases. There are four isoforms of TNNI3K that are produced as a result of alternative splicing events.	Isotype: IgG SWISS: Q59H18	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200) Reactivity: Human, Rat (predicted: Mouse, Rabbit, Sheep, Cow, Chicken, Horse) Predicted MW.: 93 kDa Subcellular Location: Cytoplasm ,Nucleus
---	---	---

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TNNI3K) Polyclonal Antibody, Unconjugated (bs-9458R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TNNI3K) Polyclonal Antibody, Unconjugated (bs-9458R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.