

bs-20166R**[Primary Antibody]****HCN2 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 610 Target: HCN2 Immunogen: KLH conjugated synthetic peptide derived from human HCN2: 651-750/889. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Hyperpolarization-activated, cyclic nucleotide-binding channels (HCN) are voltage-gated cation channels that are activated by direct binding of intracellular cyclic nucleotides. The HCN family consists of four members (HCN1–4), each with a core transmembrane segment domain and a carboxy-terminal 120 amino-acid cyclic nucleotide-binding domain motif (1). HCN channels are expressed in the brain, heart, thalamus and testis (1). The pacemaker properties of HCN channels contribute to spontaneous rhythmic activity in the brain and heart (1). The genes encoding human HCN1 and HCN2 map to chromosomes 5 and 19p13.3, respectively (2,3). The genes encoding HCN3 and HCN4 map to chromosomes 1q21.3 and 15q24-q25, respectively (4,5).	Isotype: IgG SWISS: Q9UL51 Applications: WB (1:500-2000) Reactivity: Human (predicted: Mouse, Rat, Rabbit, Sheep, Cow, Dog) Predicted MW.: 97 kDa Subcellular Location: Cell membrane
--	--

— VALIDATION IMAGES —

Sample: Lane 1: Human Jurkat cell lysates Lane 2: Human SH-SY5Y cell lysates Primary: Anti-HCN2 (bs-20166R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 97 kDa Observed band size: 110 kDa

— SELECTED CITATIONS —

- **[IF=1.468]** Fei-Fei Wang, et al. Aging-induced atrial fibrosis in If current change and its effect on atrial fibrillation in dogs. ANN NONINVAS ELECTRO. 2022 Apr 11 WB ;Dog. 35403309