bs-20167R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

HCN2 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 610 SWISS: Q9UL51

Target: HCN2

Immunogen: KLH conjugated synthetic peptide derived from human HCN2:

61-150/889.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS,

pH7.4.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Hyperpolarization-activated, cyclic nucleotide-binding channels

(HCN) are voltage-gated cation channels that are activated by direct binding of intracellular cyclic nucleotides. The HCN family

consists of four members (HCN1–4), each with a core

transmembrane segment domain and a carboxy-terminal 120 amino-acid cyclic nucleotide-binding domain motif (1). HCN

channels are expressed in the brain, heart, thalamus and testis (1). The pacemaker properties of HCN channels contribute to spontaneous rhythmic activity in the brain and heart (1). The genes

spontaneous rhythmic activity in the brain and heart (1). The genes encoding human HCN1 and HCN2 map to chromosomes 5 and 19p13.3, respectively (2,3). The genes encoding HCN3 and HCN4 map to chromosomes 1q21.3 and 15q24-q25, respectively (4,5).

Applications: IHC-P (1:100-500)

IHC-F (1:100-500)


IF (1:100-500)

Reactivity: Human

Predicted MW.: 97 kD

Subcellular Location: Cell membrane

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Human cervical cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HCN2) Polyclonal Antibody, Unconjugated (bs-20167R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.