

bs-19536R

[Primary Antibody]

ZNF502 Rabbit pAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 91392	SWISS: Q8TBZ5	
Target: ZNF502		Reactivity: Human
Immunogen: KLH conjugated synthetic peptide derived from human ZNF502: 5-150/544.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 63 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Nucleus
Background: ZNF502 may be involved in transcriptional regulation. It belongs to the krueppel C2H2 type zinc finger protein family. ZNF502 contains fourteen C2H2 type zinc fingers.		

— VALIDATION IMAGES —



Sample: Lane 1: Human SH-SY5Y cell lysates
Lane 2: Human U-2 OS cell lysates Lane 3:
Human HUVEC cell lysates Lane 4: Human 293T
cell lysates Lane 5: Human A549 cell lysates
Primary: Anti-ZNF502 (bs-19536R) at 1/1000
dilution Secondary: IRDye800CW Goat Anti-
Rabbit IgG at 1/20000 dilution Predicted band
size: 63 kDa Observed band size: 60 kDa



Tissue/cell: human brain glioma; 4%
Paraformaldehyde-fixed and paraffin-
embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block
endogenous peroxidase by 3% Hydrogen
peroxide for 30min; Blocking buffer (normal goat
serum,C-0005) at 37℃ for 20 min; Incubation:
Anti-ZNF502 Polyclonal Antibody,
Unconjugated(bs-19536R) 1:500, overnight at
4℃, followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining