

ZNF502 Rabbit pAb

Catalog Number: bs-19536R

Target Protein: ZNF502

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human

Predicted MW: 63 kDa

Entrez Gene: 91392

Swiss Prot: Q8TBZ5

Source: KLH conjugated synthetic peptide derived from human ZNF502: 5-150/544.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: ZNF502 may be involved in transcriptional regulation. It belongs to the krueppel C2H2 type zinc finger protein family. ZNF502 contains fourteen C2H2 type zinc fingers.

VALIDATION IMAGES



Sample: Lane 1: Human SH-SY5Y cell lysates Lane 2: Human U-2 OS cell lysates Lane 3: Human HUVEC cell lysates Lane 4: Human 293T cell lysates Lane 5: Human A549 cell lysates Primary: Anti-ZNF502 (bs-19536R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 63 kDa Observed band size: 60 kDa



Tissue/cell: human brain glioma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-ZNF502 Polyclonal Antibody, Unconjugated(bs-19536R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining