

bs-13626R**[Primary Antibody]****CED6 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse (predicted: Rabbit, Pig, Sheep, Cow, Horse) Predicted MW.: 34 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 51454	SWISS: Q9UBP9	
Target: CED6		
Immunogen: KLH conjugated synthetic peptide derived from human CED6: 31-130/304.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Several proteins involved in regulating and executing programmed cell death have been identified in <i>C. elegans</i> . CED-3, a member of the ICE protease/caspase family, and CED-4, a homolog of the mammalian Apaf-1, promote apoptosis. CED-9, a homolog of the mammalian Bcl-2 protein, inhibits cell death. EGL-1 and CED-6 both function as death-promoting proteins, with CED-6 playing a role in the engulfment of apoptotic cells. CED-5 and CED-7 are <i>C.</i> <i>elegans</i> orthologs of the mammalian DOCK180 and ABC transporter proteins, respectively, and also play a role in the engulfment of dying cells.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CED6) Polyclonal Antibody, Unconjugated (bs-13626R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CED6) Polyclonal Antibody, Unconjugated (bs-13626R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.