
DIP13B Rabbit pAb

Catalog Number: bs-13631R

Target Protein: DIP13B

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Rat (predicted: Mouse, Sheep, Cow, Chicken, Dog)

Predicted MW: 74 kDa

Entrez Gene: 55198

Swiss Prot: Q8NEU8

Source: KLH conjugated synthetic peptide derived from human DIP13B/APPL2: 101-200/664.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The APPL family of proteins are involved in linking, trafficking and signaling downstream of tyrosine kinase receptors. APPL1, also designated adaptor protein containing pH domain, PTB domain and leucine zipper motif 1; APPL; or DCC interacting protein 13Å (DIP13Å), and APPL2, also designated adaptor protein containing pH domain, PTB domain and leucine zipper motif 2 or DCC interacting protein 13 Å (DIP13 Å), are involved in the coupling of epidermal growth factor (EGF) signaling and chromatin remodeling in the nucleus. They associate with GTPase Rab 5 and are released from the plasma membrane and translocated to the nucleus. In the nucleus, APPL1 and APPL2 associate with NuRD/MeCP1 and are essential for cell growth and proliferation. APPL2 also associates with follicle stimulating hormone receptor (FSHR). APPL2 is highly expressed in heart, brain, skeletal muscle, and kidney. APPL2 shares 54% homology with APPL1


VALIDATION IMAGES



Sample:Lovo Cell (Human) Lysate at 30 ug Primary: Anti-DIP13B (bs-13631R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 74kD Observed band size: 76kD



Sample:293T Cell (Human) Lysate at 30 ug Primary: Anti-DIP13B (bs-13631R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 74kD Observed band size: 76kD



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DCC interacting protein 13 beta; DIP13B) Polyclonal Antibody, Unconjugated (bs-13631R) at 1:200 overnight at 4°C, followed by a conjugated secondary antibody (bs-0295G-cy3) for 90 minutes .