
EMR2 Rabbit pAb

Catalog Number: bs-13634R

Target Protein: EMR2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human, Mouse

Predicted MW: 88 kDa

Entrez Gene: 30817

Swiss Prot: Q9UHX3

Source: KLH conjugated synthetic peptide derived from human EMR2: 451-550/823.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The epidermal growth factor (EGF) family constitutes a group of class B, G protein-coupled receptors, which includes CD97 and EMR2. EMR2 is a member of the EGF-TM7 receptor subfamily. EGF-TM7 receptors are a family of class B, seven-span transmembrane (TM7) receptors predominantly expressed by cells of the immune system. Within the TM7 superfamily, the molecular structure and ligand-binding properties of EGF-TM7 receptors are unique. Derived from the processing of a single polypeptide, they are expressed at the cell surface as heterodimers consisting of a large extracellular region associated with a TM7 moiety. Through a variable number of N-terminal EGF-like domains, EGF-TM7 receptors interact with cellular ligands such as CD55 and chondroitin sulfate. EMR2 is a heptahelical molecule predominantly expressed on cells of the immune system such as leukocytes. EMR2 is proteolytically cleaved into two separate subunits: a seven-transmembrane subunit, and an extracellular Alpha subunit.

VALIDATION IMAGES



Sample: TM4(Mouse) Cell Lysate at 30 ug MCF-7(Human) Cell Lysate at 30 ug Primary: Anti-EMR2 (bs-13634R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 88 kD Observed band size: 128 kD