
Recombinant H5N6-NA protein, His

Catalog Number: bs-49058P

Concentration: >0.5 mg/ml

AA Seq: 2-459/459

Predicted MW: 50

Detected MW: 55 kDa

Tags: His

Activity: Not tested

Endotoxin: Not analyzed

Purity: >95% as determined by SDS-PAGE

Purification: AC

Form: Lyophilized or Liquid

Storage: 20mM Tris-HCl (pH8.0) with 150mM NaCl.

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: Catalyzes the removal of terminal sialic acid residues from viral and cellular glycoconjugates. Cleaves off the terminal sialic acids on the glycosylated HA during virus budding to facilitate virus release. Additionally helps virus spread through the circulation by further removing sialic acids from the cell surface. These cleavages prevent self-aggregation and ensure the efficient spread of the progeny virus from cell to cell. Otherwise, infection would be limited to one round of replication. Described as a receptor-destroying enzyme because it cleaves a terminal sialic acid from the cellular receptors. May facilitate viral invasion of the upper airways by cleaving the sialic acid moieties on the mucin of the airway epithelial cells. Likely to play a role in the budding process through its association with lipid rafts during intracellular transport. May additionally display a raft-association independent effect on budding. Plays a role in the determination of host range restriction on replication and virulence. Sialidase activity in late endosome/lysosome traffic seems to enhance virus replication.

VALIDATION IMAGES



The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.