
Recombinant human TNF-Alpha protein (Active, HEK293)

Catalog Number: bs-48004P

Concentration: >0.5 mg/ml

AA Seq: 77-233/233

Predicted MW: 17

Detected MW: 17 kDa

Tags: Tag free

Activity: Yes

Endotoxin: <0.5 EU/μg as determined by LAL.

Purity: >95% as determined by SDS-PAGE

Purification: AC


Form: Lyophilized

Storage: Lyophilized from 0.22μm filtered solution in PBS (pH7.4) with 5mM DTT. Normally 5% trehalose is added as protectant before Lyophilization.

Use a manual defrost freezer and avoid repeated freeze-thaw cycles. In general: 12 months from date of receipt, -20 to -80°C as supplied. 1 month, 2 to 8°C under sterile conditions after reconstitution. 3 months, -20 to -80°C under sterile conditions after reconstitution.

Background: This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, psoriasis, rheumatoid arthritis ankylosing spondylitis, tuberculosis, autosomal dominant polycystic kidney disease, and cancer. Mutations in this gene affect susceptibility to cerebral malaria, septic shock, and Alzheimer disease. Knockout studies in mice also suggested the neuroprotective function of this cytokine. [provided by RefSeq, Aug 2020]

VALIDATION IMAGES



The purity of the protein is greater than 95% as determined by non reduced and reduced SDS-PAGE.