

bs-0104M**[Primary Antibody]****beta Amyloid 1-16 Mouse pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Mouse	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 351	SWISS: P05067	IHC-F (1:100-500)
Target: beta Amyloid 1-16		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide of human beta-Amyloid: 1-16/42.		Reactivity: Human, Rat, Rabbit (predicted: Pig)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 4.3/83 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cell membrane
Background: The cerebral and vascular plaques associated with Alzheimer's disease are mainly composed of Amyloid beta peptides. beta Amyloid is derived from cleavage of the Amyloid precursor protein and varies in length from 39 to 43 amino acids. beta Amyloid [1-40], beta Amyloid [1-42], and beta Amyloid [1-43] peptides result from cleavage of Amyloid precursor protein after residues 40, 42, and 43, respectively. The cleavage takes place by gamma-secretase during the last Amyloid precursor protein processing step. beta Amyloid [1-40], beta Amyloid [1-42], and beta Amyloid [1-43] peptides are major constituents of the plaques and tangles that occur in Alzheimer's disease. beta Amyloid antibodies and peptides have been developed as tools for elucidating the biology of Alzheimer's disease.		

— VALIDATION IMAGES —

Sample: Lane 1: Human U87MG Cell Lysates
 Lane 2: Human SY5Y cell lysates Lane 3: Human U251 Cell Lysates
 Primary: Anti-beta Amyloid 1-16 (bs-0104M) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution
 Predicted band size: 4.3/83 kDa
 Observed band size: 130 kDa



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (beta Amyloid 1-16) Monoclonal Antibody, Unconjugated (bs-0104M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.