

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Protein A / FITC

Catalog Number: bsF-0362P Concentration: 1mg/ml

AA Seq: Purified native protein

Activity: Not tested Endotoxin: Not analyzed

Form: Liquid

Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: Protein A is a 40-60 kDa surface protein originally found in the cell wall of the bacteria

notably IgGs, and helps inhibit phagocytic engulfment and acts as an immunological disguise via this type of interaction, thus the bacteria will disrupts opsonization and phagocytosis. SPA is known to bind with Fc region of immunoglobulins preferentially through interaction with the VH3 variable region of the heavy chain. SPA has been shown to

Staphylococcus aureus (SPA). SPA binds proteins from many of mammalian species, most

bind with high affinity to human IgG1 and IgG2 as well as mouse IgG2a and IgG2b, whereas bind with moderate affinity to human IgM, IgA and IgE as well as mouse IgG3 and IgG1. SPA is often produced in E. coli and is practically coupled to other molecules such as enzymes, biotin, radioactive iodine for use in immunology and other biological research. SPA is also immobilized onto solid supports such as agarose beads for total IgG purifying or interest

protein or protein complex identifying in immunoprecipitation studies.